SATEJ
GLOBAL SCIENCE
Think Globally, Act Locally

Plot No. 272/A, Opp. Renaissance Hotel,
Chankyapuri Road, Nr. Kargil Petrol Pump,
Off. S.G. Highway, Ahmedabad- 380061.
+91 9723287125
info@satejglobalscience.com
www.satejglobalscience.com

Myoma


Overview

Myoma or uterine fibroids are benign tumors of muscle and connective tissue that develop within or are attached to the uterine wall.

The cause of fibroid tumors of the uterus is unknown. However, it is suggested that fibroids may enlarge with estrogen therapy (such as oral contraceptives) or with pregnancy.

Their growth seems to depend on regular estrogen stimulation, showing up only rarely before the age of 20 and shrinking after menopause. As long as a woman with fibroids is menstruating, the fibroids will probably continue to grow, although growth is usually quite slow.

Fibroids can be microscopic, but they can also grow to fill the uterine cavity, and may weigh several pounds. Uterine fibroids are the most common pelvic tumor and they may be present in 15 to 20% of reproductive-age women, and 30 to 40% of women over 30.



A pelvic examination reveals an irregularity shaped, lumpy or enlarged uterus. Frequently, this diagnosis is reliable, however, on occasion; diagnosis is difficult, especially in obese women. Fibroid tumors have been mistaken for ovarian tumors. Inflammatory processes of the tubes, and pregnancy

It is possible for a single fibroid to develop although normally there are a number of them, which begin as small seedlings spread throughout the muscular walls of the uterus.

They slowly enlarge and become more nodular, frequently intruding into the cavity of the uterus or growing out beyond the normal boundary of the uterus. Rarely, a fibroid will hang from a long stalk attached to the outside of the uterus. This is called a pedunculated fibroid

SATEJ
GLOBAL SCIENCE
Think Globally, Act Locally

Plot No. 272/A, Opp. Renaissance Hotel,
Chankyapuri Road, Nr. Kargil Petrol Pump,
Off. S.G. Highway, Ahmedabad- 380061.
+91 9723287125
info@satejglobalscience.com
www.satejglobalscience.com

Signs and Symptoms

- Sensation of fullness or pressure in lower abdomen
- Abdominal fullness, gaseous
- Pelvic cramping or pain with periods
- Increase in urinary frequency
- Heavy menstrual bleeding (menorrhagia), sometimes with the passage of blood clots, sometimes resulting in anemia (a low blood count)
- Sudden, severe pain (a pedunculated fibroid twists causing the blood vessels feeding the tumor to kink) hospitalization and surgery may be needed in this instance.

Benefit of SGS Therapy:

- Help in removing uterine Fibroids
- Regulate the general metabolism.
- Below combined supplement therapy gives very synergetic effects as an anti-tumours & anti Systs.
- Make normal cell mitosis functioning & cellular metabolism
- Provides good hormonal balance in women and ensures Minimum miscarriages
- Decisively Provide healthy Nourishment in women's reproductive system & healthy fetus

Suggested Products:

Minerjal : Take 10 drops 3 times in 100 ml of Drinking water
Minerjal Lyte : Add 25 drops in 1 litre Drinking water and finishes it in a day
Grolyfe : Take 10 drops 3 times under the tongue
Comraid : 2 Capsules 3 times after meal
LivoSat : Daily 1 Sachet 2 time









Copyright © SGS